

# Optimal Execution under Liquidity Uncertainty

Sergio Pulido



Joint work with Etienne Chevalier, Yadh Hafsi and Vathana Ly Vath

Quantitative Finance at La Défense

June 18, 2026

# Introduction

## The Execution Problem:

- An agent must execute  $\bar{X}$  number of shares of a financial asset by time  $T < +\infty$ .
- The agent is a trader with continuous observations of the Limit Order Book (LOB).
- Both continuous and discrete trading are allowed.

## Optimal Execution with transient price impact:

- Obizhaeva and Wang (2013), Alfonsi, Fruth and Schied (2010), Predoiu, Shaikhet, and Shreve (2011), Gatheral, Schied and Slynko (2012), Fruth, Schöneborn and Urusov (2014), Alfonsi and Blanc (2016), Abi Jaber and Neuman (2025) ...

## Optimal Execution with stochastic liquidity/resilience:

- Bayraktar and Ludkovski (2011), Becherer, Bilarev and Frentrup (2018), Fruth, Schoneborn, and Urusov (2019), Ackermann, Kruse, and Urusov (2021), Fouque, Jaimungal, and Saporito (2022), Cartea, Sanchez-Betancourt (2023), Dammann and Ferrari (2023) and Chevalier, Hafsı and Ly Vath (2024) ...

## Focus:

→ Singular control problem under (stochastic) liquidity constraints.

# Introduction

## The Execution Problem:

- An agent must execute  $\bar{X}$  number of shares of a financial asset by time  $T < +\infty$ .
- The agent is a trader with continuous observations of the Limit Order Book (LOB).
- Both continuous and discrete trading are allowed.

## Optimal Execution with transient price Impact:

- Obizhaeva and Wang (2013), Alfonsi, Fruth and Schied (2010), Predoiu, Shaikhet, and Shreve (2011), Gatheral, Schied and Slynko (2012), Fruth, Schöneborn and Urusov (2014), Alfonsi and Blanc (2016), Abi Jaber and Neuman (2025) ...

## Optimal Execution with stochastic liquidity/resilience:

- Bayraktar and Ludkovski (2011), Becherer, Bilarev and Frentrup (2018), Fruth, Schoneborn, and Urusov (2019), Ackermann, Kruse, and Urusov (2021), Fouque, Jaimungal, and Saporito (2022), Cartea, Sanchez-Betancourt (2023), Dammann and Ferrari (2023) and Chevalier, Hafsi and Ly Vath (2024) ...

## Focus:

→ Singular control problem under (stochastic) liquidity constraints.

# Introduction

## The Execution Problem:

- An agent must execute  $\bar{X}$  number of shares of a financial asset by time  $T < +\infty$ .
- The agent is a trader with continuous observations of the Limit Order Book (LOB).
- Both continuous and discrete trading are allowed.

## Optimal Execution with transient price Impact:

- Obizhaeva and Wang (2013), Alfonsi, Fruth and Schied (2010), Predoiu, Shaikhet, and Shreve (2011), Gatheral, Schied and Slynko (2012), Fruth, Schöneborn and Urusov (2014), Alfonsi and Blanc (2016), Abi Jaber and Neuman (2025) ...

## Optimal Execution with stochastic liquidity/resilience:

- Bayraktar and Ludkovski (2011), Becherer, Bilarev and Frentrup (2018), Fruth, Schoneborn, and Urusov (2019), Ackermann, Kruse, and Urusov (2021), Fouque, Jaimungal, and Saporito (2022), Cartea, Sanchez-Betancourt (2023), Dammann and Ferrari (2023) and Chevalier, Hafsi and Ly Vath (2024) ...

## Focus:

→ Singular control problem under (stochastic) liquidity constraints.

# Introduction

## The Execution Problem:

- An agent must execute  $\bar{X}$  number of shares of a financial asset by time  $T < +\infty$ .
- The agent is a trader with continuous observations of the Limit Order Book (LOB).
- Both continuous and discrete trading are allowed.

## Optimal Execution with transient price Impact:

- Obizhaeva and Wang (2013), Alfonsi, Fruth and Schied (2010), Predoiu, Shaikhet, and Shreve (2011), Gatheral, Schied and Slynko (2012), Fruth, Schöneborn and Urusov (2014), Alfonsi and Blanc (2016), Abi Jaber and Neuman (2025) ...

## Optimal Execution with stochastic liquidity/resilience:

- Bayraktar and Ludkovski (2011), Becherer, Bilarev and Frentrup (2018), Fruth, Schoneborn, and Urusov (2019), Ackermann, Kruse, and Urusov (2021), Fouque, Jaimungal, and Saporito (2022), Cartea, Sanchez-Betancourt (2023), Dammann and Ferrari (2023) and Chevalier, Hafsi and Ly Vath (2024) ...

## Focus:

→ Singular control problem under (stochastic) liquidity constraints.

# Liquidity Dynamics: Market Impact

- Fix a liquidity regime  $i$
- Impact function derived from the limit order book (LOB) shape  $F_i$ .
- Quantity at distance  $x \geq 0$  from fundamental price  $A_t$  for a continuous LOB:  
$$F_i(x) := \int_0^x f_i(p) dp.$$
- Price impact for order size  $y$ :  $\psi_i(y) := \sup\{a \geq 0 : F_i(a) < y\}$ , with  $\psi_i(0) = 0$ .
- Post-trade price:  $A_t + \psi_i(y)$ .




Figure 1: Illustration of a limit order book density and resulting price impact.

# Liquidity Dynamics: Price Modeling and Volume Effect

- **Liquidity Regimes  $I$ :** A stable and conservative Markov chain  $\{I_t\}_{t \geq 0}$  on finite state space  $E = \{1, \dots, d\}$ .
- **Volume effect  $Y$ :** an  $\mathcal{F}$ -adapted nonnegative process, such that for all  $t \geq 0$  and  $u \in [t, T]$ ,

$$\begin{cases} dY_u = -h(Y_{u-})du + \sigma(Y_{u-})dW_u + \int_{\mathbb{R}} q(Y_{u-}, z)M(du, dz), \text{ (comp. of } M(du, dz) = \lambda_u \nu(dz)du) \\ Y_{t-} = y. \end{cases}$$

- **Price  $P$ :**  $P_t := \underbrace{A_t}_{\text{fundamental price}} + \underbrace{D_t}_{\text{price deviation}}, \text{ with } D_t := \psi_t(Y_t), \text{ for all } t \geq 0.$




Figure 2: Representation of a limit order book at time  $t$ . The left-hand figure showing the state before a transaction and the right-hand figure showing it after. The dashed area  $Y_t$  denotes consumed shares.

# Problem Formulation

## Objective:

→ Purchase a position  $\bar{X}$  within a finite time horizon  $T$ , starting from  $X_{t-} = x$  if  $I_{t-} = i$  and  $Y_{t-} = y$ .

- Supposing  $y = 0$ , the purchase cost for a large trade of size  $\Delta y$ , in excess of  $A_t$  at time  $t$  is equal to

$$\Phi_i(\Delta y) := \int_0^{\psi_i(\Delta y)} \xi dF_i(\xi) = \int_0^{\Delta y} \psi_i(\zeta) d\zeta, \quad \forall (y, i) \in \mathbb{R}_+ \times \mathbb{I}_m.$$

- The total cost is

$$\pi_t^i(\Delta y) = \int_0^{\Delta y} [A_t + \psi_i(\zeta)] d\zeta = \underbrace{A_t \Delta y}_{\text{Cost at the current price}} + \underbrace{\Phi_i(\Delta y)}_{\text{Impact cost}}.$$

- An admissible purchase strategy consists of a non-decreasing  $\mathcal{F}$ -adapted right-continuous process  $X = (X_u)_{t \leq u \leq T}$  such that  $X_{t-} = x$  and  $X_T = \bar{X}$ .

# Optimal Execution: Value Function

## Controlled Dynamics:

$$\begin{cases} dY_u^{t,y,X} = dX_u - h(Y_{u^-}^{t,y,X})du + \sigma(Y_{u^-}^{t,y,X})dW_u + \int_{\mathbb{R}} q(Y_{u^-}^{t,y,X}, z)M(du, dz), \\ Y_t^{t,y,X} = y, \end{cases}$$

with  $u \in [t, T]$  and  $y \geq 0$ .

## Value Function:

- Minimize the costs over admissible strategies  $X \in \mathcal{A}_t(x)$ :

$$v(i, t, x, y) = v_i(t, x, y) := \inf_{X \in \mathcal{A}_t(x)} \mathbb{E} \left[ \int_t^T \psi_{l_u}(\check{Y}_{u^-}^{t,y,X}) dX_u^c + \sum_{t \leq u \leq T} \Phi_{l_u}(Y_u^{t,y,X}) - \Phi_{l_u}(\check{Y}_{u^-}^{t,y,X}) \right],$$

for all  $(t, x, y) \in \mathcal{S} := [0, T] \times [0, \bar{X}] \times \mathbb{R}_+^*$  and  $i \in \mathbb{I}_m$ .

→ *Boundary conditions:*  $v_i(T, x, y) = \Phi_i(y + \bar{X} - x) - \Phi_i(y)$  and  $v_i(t, \bar{X}, y) = 0$ .

→ *Growth condition:*  $0 \leq v_i(t, x, y) \leq \Phi_i(y + \bar{X} - x) - \Phi_i(y)$ .

# Optimal Execution: Dynamic Programming Principle and HJBQVI

## Value Function and HJBQVI:

- $v$  solves the Hamilton-Jacobi-Bellman Quasi-Variational Inequality (HJBQVI):

$$\max \left( -\frac{\partial v_i}{\partial t} - \mathcal{L} v_i - \sum_{j \neq i} (v_j - v_i) Q_{ij}, -\frac{\partial v_i}{\partial x} - \frac{\partial v_i}{\partial y} - \psi_i \right) = 0, \text{ on } \mathcal{S}.$$

where  $\mathcal{S} := [0, T] \times [0, \bar{X}] \times \mathbb{R}_+$ .  $\rightarrow$  **viscosity solutions**

- The partial integro-differential operator  $\mathcal{L}$  is given by

$$\mathcal{L} \varphi := \frac{1}{2} \sigma^2(y) \frac{\partial^2 \varphi}{\partial y^2} - h(y) \frac{\partial \varphi}{\partial y} + \lambda_t \int_{\mathbb{R}} (\varphi(t, x, y + q(y, z)) - \varphi) v(dz),$$

## Theorem 1 (Dynamic Programming Principle and HJBQVI)

The value function  $v$  is a continuous function on  $\bar{\mathcal{S}}$ . For any stopping time  $\tau$  in  $[t, T]$ ,  $(t, x, y) \in \mathcal{S}$  and  $i \in \mathbb{I}_m$ , we have

$$v_i(t, x, y) = \inf_{X \in \mathcal{A}_t^c(x)} \mathbb{E} \left[ \int_t^\tau \psi_{l_{u^-}}(\check{Y}_{u^-}^{t,y,X}) dX_u^c + \sum_{t \leq u \leq \tau} (\Phi_{l_u}(Y_u^{t,y,X}) - \Phi_{l_u}(\check{Y}_{u^-}^{t,y,X})) + v_{l_\tau}(\tau, X_\tau, Y_\tau^{t,y,X}) \right].$$

The value function  $v$  is the unique viscosity solution of the HJBQVI.

# The Free Boundary Problem

## Definition 3.1

We define the exercise region  $\mathcal{E}_i := \overline{\text{int}(\mathcal{E}_i^{\text{diff}})}$  as the closure of interior of the set  $\mathcal{E}_i^{\text{diff}}$ , on which  $v_i$  is differentiable, with

$$\mathcal{E}_i^{\text{diff}} := \left\{ (t, x, y) \in \mathcal{S} \setminus \mathcal{N}_i : -\frac{\partial v_i}{\partial x} - \frac{\partial v_i}{\partial y} - \psi_i = 0 \right\},$$


and we define the continuation region  $\mathcal{C}_i := \mathcal{S} \setminus \mathcal{E}_i$  as its complement.

## Proposition 3.1 (Connectedness)

Assume that the interiors of  $\mathcal{C}_i$  and  $\mathcal{E}_i$  are non-empty, and that  $\sigma + \lambda > 0$ . Then, for each  $i \in \mathbb{I}_m$ , the free boundary  $\partial \mathcal{E}_i$  is non-empty and path-connected. Moreover, if  $(t, x, y) \in \mathcal{E}_i$ , then

$$(t, x, y') \in \mathcal{E}_i \text{ for all } 0 \leq y' \leq y, \text{ and } (t, x', 0) \in \mathcal{E}_i \text{ for all } 0 \leq x' \leq x.$$

## Illustration of the boundary



**Figure 3:** Illustration of continuation (blue) and exercise (red) regions, plotted against purchased quantity (x-axis) and volume effect (y-axis). The arrow shows an impulse shifting a state from exercise to continuation along  $y = x$ .

## Numerical Examples: Parameterization

- Volume effect:

$$h(y) = cy, \quad \sigma(y) = dy, \quad \text{and} \quad q(y, z) = eyz, \quad \forall y, z \in \mathbb{R}_+,$$

- Order volumes follow an exponential distribution:  $\nu(dz) = \eta e^{-\eta z} \mathbb{1}_{\{z>0\}} dz, \quad \forall z \in \mathbb{R}.$
- Limit order book shape and impact functions:

$$\psi_i(y) = \begin{cases} \frac{y}{\kappa}, & \text{if } \gamma_i = 0, \\ e^{\frac{y}{\kappa}} - 1, & \text{if } \gamma_i = 1, \\ [1 + (1 - \gamma_i) \frac{y}{\kappa}]^{\frac{1}{1-\gamma_i}} - 1, & \text{else.} \end{cases}$$

- Default parameter values:

Parameter	$c$	$d$	$e$	$\eta$	$\kappa$	$\gamma_0$	$\lambda$	$\bar{X}$	$\bar{Y}$	$T$
Value	0.5	0.1	0.2	1.0	0.8	-1	0.5	4	10	4.0

## Numerical Results: Single Regime Case

→ Impact of drift  $h$ , jump intensity  $\lambda$ , and volatility  $\sigma$  on the exercise and continuation regions.




Figure 4: Variation of the exercise boundary  $x \mapsto y^*(T/2, x)$  under different market conditions with jump intensity on the left, resilience in the middle, and volatility on the right.

## Numerical Results: Regime-Switching case

We assume a two-state homogeneous Markov chain, with the transition rate matrix  $Q(t)$  at time  $t \in [0, T]$  given by

$$Q(t) = \begin{pmatrix} -q_1 & q_1 \\ q_2 & -q_2 \end{pmatrix}.$$

We fix  $q_1 = q_2 = 0.2$  and  $\gamma_1 = 0$ . With these parameters set, we then vary  $\gamma_2$ .




Figure 5: Variation of the exercise boundary  $x \mapsto y^*(T/2, x)$  of  $v_0$  on the left and  $v_1$  on the right under different price impacts.

# Conclusions


- Developed a stochastic control framework for optimal execution under general price impact and stochastic liquidity, modeled via a jump diffusion volume effect and regime-switching Markov processes.
- Characterized the value function as the unique viscosity solution of a system of HJBQVIs. Proved regularity properties of the value function, including continuity, monotonicity, and connectedness of the free boundary.
- Approximated the solution to quantify the effect of liquidity uncertainty on optimal strategies.

# Thank You !

Chevalier, E., Hafsi, Y., Ly Vath, V., Pulido, S. (2026). Optimal Execution under Liquidity Uncertainty. arXiv:2506.11813. To appear in SIAM Journal on Financial Mathematics.

## Numerical Results: Single Regime Case

→ Impact of drift  $h$ , jump intensity  $\lambda$ , and volatility  $\sigma$  on the execution costs (i.e, value function).



**Figure 6:** Variation of the value function over time under different market conditions with jump intensity on the left, resilience in the middle, and volatility on the right.

## Numerical Results: Regime-Switching case

→ The price impact parameter is set to  $\gamma_1 = 0$  in regime 1 and  $\gamma_2 = 0.5$  in regime 2. We consider  $q_1 = q_2 = q_0$  and vary the values of  $q_0$ . The probability of being in either regime is equal, but the frequency of transitions between regimes changes.




Figure 7: Variation of the exercise boundary  $x \mapsto y^*(T/2, x)$  of  $v_0$  on the left and  $v_1$  on the right under different regime switching intensities.

## Numerical Results: Regime-Switching case

→ The price impact parameter is set to  $\gamma_1 = 0$  in regime 1 and  $\gamma_2 = 0.5$  in regime 2. We fix  $q_1 = 0.2$  and vary  $q_2$ , and study how the asymmetry in switching probabilities affects execution.




Figure 8: Variation of the exercise boundary  $x \mapsto y^*(T/2, x)$  of  $v_0$  on the left and  $v_1$  on the right under different asymmetric regime switching intensities.

→ Stochastic liquidity and regime changes.




Figure 9: Changes in the spread observed in BNP stock data on 11/05/2022, Chevalier, Hafsi, and Ly Vath (2023).

# Liquidity Dynamics: Price Modeling and Volume Effect

## Assumptions 4.1

(A1) *There exists  $b > 0$ ,  $\beta > 0$  and  $a > 0$  such that*

$$F_i(x) \geq bx^\beta, \quad \forall (x, i) \in [a, +\infty[ \times \mathbb{I}_m.$$

(A2) *The measure  $\nu(dz)$  satisfies*

$$\int_{\mathbb{R}} (1 + z^2) \nu(dz) < +\infty,$$

*and  $y \mapsto y + q(y, z)$  is non-decreasing for every  $z$*

(A3) *There exists a constant  $C > 0$  such that, for all  $y \geq 0$ ,*

$$|h(y)| + |\sigma(y)| + \int_{\mathbb{R}} |q(y, z)| \nu(dz) \leq C(1 + |y|).$$

(A4) *There exists a constant  $L > 0$  such that, for all  $y, y' \geq 0$ ,*

$$|h(y) - h(y')| + |\sigma(y) - \sigma(y')| + \int_{\mathbb{R}} |q(y, z) - q(y', z)| \nu(dz) \leq L|y - y'|.$$

# Analytical Properties of the Value Function

## Proposition 4.1

*The value function  $v$  is finite.*

## Proposition 4.2 (Monotonicity)

*For any  $(i, t, x, y) \in \mathbb{I}_m \times \overline{\mathcal{F}}$ , the following results hold:*


- 1  $t_0 \mapsto v_i(t_0, x, y)$  is non-decreasing on  $[0, T]$ .
- 2  $x_0 \mapsto v_i(t, x_0, y)$  is non-increasing on  $[0, \bar{X}]$ .
- 3  $y_0 \mapsto v_i(t, x, y_0)$  is non-decreasing on  $\mathbb{R}_+$ .

## Theorem 4.1

*The value function  $v$  is a continuous function on  $\overline{\mathcal{F}}$ .*

## Analytical Properties of the Value Function

**Sketch of proof:** Compare the sample paths of  $Y^{t,y,X}$  and  $Y^{t,y',\tilde{X}}$ . Design a strategy that ensures the paths of  $u \mapsto Y_u^{t,y,X}$  and  $u \mapsto Y_u^{t,y',\tilde{X}}$  meet at a specific point in time  $\tau$ , facilitating their comparison and enabling us to establish bounds for  $v_i(t,x,y') - v_i(t,x,y)$ .



**Figure 10:** Illustration of  $Y^{t,y,X}$  and  $Y^{t,y',\tilde{X}}$  sample paths in the scenario where  $y < y'$ ,  $\tau < \theta < T$  and  $\Delta X_\tau + Y_\tau^{t,y,X} - Y_\tau^{t,y',\tilde{X}} < \bar{X} - \hat{X}_{\tau-}$ .

# Viscosity Characterization

## Definition 4.1 (Viscosity Solution)

We define a viscosity solution of HJBQVI as follows :

- ①  $v$  is a continuous viscosity supersolution (resp. subsolution) of HJBQVI on  $\mathbb{I}_m \times \mathcal{S}$  if it satisfies the growth conditions, and if

$$\max \left( - \left( \frac{\partial \varphi}{\partial t} + \mathcal{L} \varphi + \sum_{j \neq i} (v_j - v_i) Q_{ij} \right) (t, x, y), - \left( \frac{\partial \varphi}{\partial x} + \frac{\partial \varphi}{\partial y} + \psi_i \right) (t, x, y) \right) \geq (\text{resp. } \leq) 0,$$

for any  $(i, t, x, y) \in \mathbb{I}_m \times \mathcal{S}$  and any smooth test function  $\varphi \in C^{1,2}(\mathcal{S})$  such that  $(v_i - \varphi)$  attains a local minimum (resp. maximum) at  $(t, x, y)$  over the set

$[t, t + \delta] \times [x, x + \delta] \times B_\delta(y) \subset \mathcal{S}$  for some  $\delta > 0$ , with  $(v_i - \varphi)(t, x, y) = 0$ .

- ②  $v$  is a continuous viscosity solution on  $\mathbb{I}_m \times \mathcal{S}$  if it is both a viscosity supersolution and subsolution of the HJBQVI.

# Viscosity Characterization

## Theorem 4.2

*The value function  $v$  is a viscosity subsolution of the HJBQVI.*

## Theorem 4.3

*The value function  $v$  is a viscosity supersolution of HJBQVI.*

## Theorem 4.4 (Strong Comparison Principle)

*If  $v_i$  is a continuous viscosity subsolution and  $w_i$  is a continuous viscosity supersolution of the HJBQVI, such that*

$$v_i(t, \bar{X}, y) \leq w_i(t, \bar{X}, y), \text{ and } v_i(T, x, y) \leq w_i(T, x, y),$$

*for all  $(i, t, x, y) \in \mathbb{I}_m \times \bar{\mathcal{S}}$ , then  $v_i \leq w_i$  on  $\mathcal{S}$ .*

## Strong comparison principle

**Sketch of proof:** Let  $\beta > 0$  such that

$$\lim_{y \rightarrow +\infty} \max_{i \in I_m} \frac{\Phi_i(y + \bar{X} - x) - \Phi_i(y)}{y^\beta} = 0, \quad \forall (x, y) \in [0, \bar{X}] \times \mathbb{R}_+.$$

Define  $\varphi_i : \bar{\mathcal{I}} \rightarrow \mathbb{R}$  such that

$$\varphi_i(t, x, y) := -e^{-ct} ((-a_1 x + a_2) y^\beta - b_1 x + b_2), \quad \forall (t, x, y) \in \bar{\mathcal{I}},$$

where  $a_1, a_2, b_1, b_2$  and  $c$  are positive constants. Define  $v_{i,m} : \bar{\mathcal{I}} \rightarrow \mathbb{R}$  such that

$$v_{i,m} := v_i + \frac{1}{m} \varphi_i.$$

$\Rightarrow v_{i,m}$  is a strict subsolution. We get a contradiction using *Ishii's lemma*.

# The Free Boundary Problem

**Sketch of proof:** Assume that  $\mathring{\mathcal{C}}_i$  and  $\mathring{\mathcal{E}}_i$  are non-empty. Let  $z_0 := (t_0, x_0, y_0) \in \mathring{\mathcal{E}}_i$  be an interior point of  $\mathcal{E}_i$ . Suppose there exists  $\delta > 0$  one of the following options,

$$\mathcal{O}_c := \{(t_0, x_0, y) \in \overline{\mathcal{F}} : y + \delta < y_0\} \subset \mathcal{E}_i, \text{ and } \mathcal{O}_e := \{(t_0, x_0, y) \in \overline{\mathcal{F}} : y_0 < y + \delta\} \subset \mathcal{E}_i,$$

$$\mathcal{O}_c := \{(t_0, x, 0) \in \overline{\mathcal{F}} : x + \delta < x_0\} \subset \mathcal{E}_i, \text{ and } \mathcal{O}_e := \{(t_0, x, 0) \in \overline{\mathcal{F}} : x_0 < x + \delta\} \subset \mathcal{E}_i,$$

where  $\mathcal{O}_c$  is non-empty segment.




Figure 11: Illustration of  $\mathcal{O}_e$ ,  $\mathcal{O}_c$  and  $\tilde{\mathcal{O}}$ .

# The Free Boundary Problem

**Sketch of proof:** Construct a test function  $\varphi_\varepsilon \in C^2(\mathcal{S})$  such that  $z_0$  achieves a local maximum of  $v_i - \varphi_\varepsilon$ ,  $v_i(z_0) = \varphi_\varepsilon(z_0)$  and  $\lim_{\varepsilon \rightarrow 0} -\mathcal{L}\varphi_\varepsilon(z_0) = +\infty$ . Since  $v_i$  is a viscosity subsolution of the HJBQVI, we get

$$\max \left( - \left( \frac{\partial \varphi_\varepsilon}{\partial t} + \mathcal{L}\varphi_\varepsilon + \sum_{j \neq i} Q_{ij}(v_j - \varphi_\varepsilon) \right) (z_0), - \left( \frac{\partial \varphi_\varepsilon}{\partial x} + \frac{\partial \varphi_\varepsilon}{\partial y} + \psi_i \right) (z_0) \right) \leq 0.$$

The constructed function  $\varphi_\varepsilon$  satisfies  $\lim_{\varepsilon \rightarrow 0} -\mathcal{L}\varphi_\varepsilon(z_0) = +\infty$ , while

$$-\left( \mathcal{L}\varphi_\varepsilon + \sum_{j \neq i} Q_{ij}(v_j - v_i) \right) (z_0) \leq 0$$

which leads to a contradiction.